

For Research Use Only

Amphiregulin Polyclonal antibody, PBS Only

Catalog Number: 16036-1-PBS

Featured Product



Basic Information

Catalog Number:

16036-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8907

GenBank Accession Number:

BC009799

GeneID (NCBI):

374

UNIPROT ID:

P15514

Full Name:

amphiregulin

Calculated MW:

252 aa, 28 kDa

Observed MW:

43 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IHC, IF-P, FC (Intra), IP, ELISA

Species Specificity:

human

Background Information

Amphiregulin (AREG) is one of the ligands of the epidermal growth factor receptor (EGFR). AREG plays a central role in mammary gland development and branching morphogenesis in organs and is expressed both in physiological and in cancerous tissues. The AREG protein is synthesized as a 252-amino acid transmembrane precursor, pro-AREG. At the plasma membrane, pro-AREG is subjected to sequential proteolytic cleavages within its ectodomain and is then released as the soluble AREG protein. Depending on the cell type and microenvironment, AREG can be produced in multiple cellular and mature forms using alternative pro-AREG cleavage sites and glycosylation motifs. Post-translational modifications of 50-kDa pro-AREG produces a major soluble 43-kDa form, 28-, 26-, 16-kDa membrane anchored forms, and soluble 21-, 19-, and 9-kDa forms (PMID: 9642297).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 16036-1-AP (Amphiregulin antibody at dilution of 1:600 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 16036-1-AP (Amphiregulin antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16036-1-AP (Amphiregulin antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 16036-1-AP (Amphiregulin antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.




IP result of anti-Amphiregulin (IP:16036-1-AP, 4ug; Detection:16036-1-AP 1:2000) with HepG2 cells lysate 3600ug. This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse embryo tissue using Amphiregulin antibody (16036-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.






1X10⁶ HeLa cells were intracellularly stained with 0.2 ug Anti-Human Amphiregulin (16036-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using Amphiregulin antibody (16036-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse embryo tissue using Amphiregulin antibody (16036-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human placenta tissue using Amphiregulin antibody (16036-1-AP) at dilution of 1:200 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Amphiregulin antibody (16036-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using Amphiregulin antibody (16036-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 16036-1-PBS in a different storage buffer formulation.