For Research Use Only

NDUFB6 Polyclonal antibody

Catalog Number:16037-1-AP

Featured Product 11 Publications




Basic Information	Catalog Number: 16037-1-AP	GenBank Accession Number: BC009801	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	600 µg/ml	4712	WB 1:1000-1:4000 IHC 1:50-1:500	
	Source: Rabbit	UNIPROT ID: 095139	IHC 1.30-1.300	
	Isotype:	Full Name:		
	IgG Immunogen Catalog Number: AG8908	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 6, 17kDa		
		Calculated MW: 128 aa, 15 kDa		
		Observed MW: 16-20 kDa		
Applications	Tested Applications:	WB, IHC, ELISA WB : A549 cells, LNCaP cells, MCF-7 cells, mouse		
	Cited Applications:			
	WB, IHC		: human liver cancer tissue,	
	Species Specificity: human, mouse, rat			
	Cited Species:			
	human, mouse			
	human, mouse Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0	atively, antigen		
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 NDUFB6, also named as NADH de amino acid protein, which belong inner membrane and is an access dehydrogenase (Complex I), that	atively, antigen d with citrate hydrogenase [ubiquinone] 1 beta sub gs to the complex I NDUFB6 subunit fa sory subunit of the mitochondrial men is believed not to be involved in cata	mily. NDUFB6 localizes in the Mitochondr nbrane respiratory chain NADH	
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 NDUFB6, also named as NADH de amino acid protein, which belong inner membrane and is an access dehydrogenase (Complex I), that electrons from NADH to the respin ubiquinone.	atively, antigen d with citrate hydrogenase [ubiquinone] 1 beta sub gs to the complex I NDUFB6 subunit fa sory subunit of the mitochondrial men is believed not to be involved in cata ratory chain. The immediate electron	lysis. Complex I functions in the transfer of acceptor for the enzyme is believed to be	
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 NDUFB6, also named as NADH de amino acid protein, which belong inner membrane and is an access dehydrogenase (Complex I), that electrons from NADH to the respin ubiquinone.	atively, antigen d with citrate hydrogenase [ubiquinone] 1 beta sub gs to the complex I NDUFB6 subunit fa sory subunit of the mitochondrial men is believed not to be involved in cata ratory chain. The immediate electron	mily. NDUFB6 localizes in the Mitochondri nbrane respiratory chain NADH Ilysis. Complex I functions in the transfer o	
Background Information	Note-IHC: suggested antige TE buffer pH 9.0; (*) Altern retrieval may be performed buffer pH 6.0 NDUFB6, also named as NADH de amino acid protein, which belong inner membrane and is an access dehydrogenase (Complex I), that electrons from NADH to the respin ubiquinone.	Antively, antigen d with citrate hydrogenase [ubiquinone] 1 beta sub gs to the complex I NDUFB6 subunit fa sory subunit of the mitochondrial men is believed not to be involved in cata ratory chain. The immediate electron Pubmed ID Journal	mily. NDUFB6 localizes in the Mitochondr nbrane respiratory chain NADH lysis. Complex I functions in the transfer of acceptor for the enzyme is believed to be Application	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16037-1-AP (NDUFB6 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 16037-1-AP (NDUFB6 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 16037-1-AP (NDUFB6 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).