For Research Use Only

## NXN Polyclonal antibody

Catalog Number:16128-1-AP

Featured Product

7 Publications




| Basic Information               | Catalog Number:<br>16128-1-AP  | GenBank Accession Number:<br>BC009327  | Purification Method:<br>Antigen affinity purification   |
|---------------------------------|--|--|---|
|                                 | Size:  | GenelD (NCBI):   | Recommended Dilutions:                                  |
|                                 | 350 µg/ml  | 64359  | WB 1:1000-1:6000  |
|                                 | Source:  | UNIPROT ID:  | IP 0.5-4.0 ug for 1.0-3.0 mg of total<br>protein lysate |
|                                 | Rabbit   | Q6DKJ4   | IHC 1:50-1:500  |
|                                 | Isotype:<br>IgG  | Full Name:<br>nucleoredoxin  | IF/ICC 1:50-1:500                                       |
|                                 | Immunogen Catalog Number:  | Calculated MW:   |   |
|                                 | AG9113   | 435 aa, 48 kDa   |   |
|                                 |  | Observed MW:<br>48 kDa   |   |
| Applications                    | Tested Applications:   | Positive   | Controls:   |
|                                 | WB, IHC, IF/ICC, IP, ELISA<br>Cited Applications:  | WB : DU 145 cells, HeLa cells, human heart tissue, HCT<br>116 cells, HEK-293 cells, HepG2 cells, SH-SY5Y cells                   |   |
|                                 | WB, IHC, IF  | IP : HeLa  | cells,  |
|                                 | Species Specificity:<br>human, mouse, rat  | IHC : hu   | man colon cancer tissue,                                |
|                                 | Cited Species:   | IF/ICC :   | HeLa cells,   |
|                                 | human, mouse, rat  |  |   |
|                                 | Note-IHC: suggested antigen retrieval with<br>TE buffer pH 9.0; (*) Alternatively, antigen<br>retrieval may be performed with citrate<br>buffer pH 6.0   |  |   |
| Background Information          | Nucleoredoxin (NRX/NXN), a relatively uncharacterized member of the TRX family protein, has recently been reported to regulate the Wnt/ $\beta$ -catenin pathway, which itself regulates cell fate and early development, in a redox dependent manner(PMID:17567240). Northern blot analysis detected several nucleoredoxin splice variants in adult mouse tissues, with highest expression in testis and skin and lowest expression in liver(PMID:9119370). It has 3 isoforms produced by alternative splicing. This antibody is specific to NXN. |  |   |
|                                 | isoforms produced by alternative   | splicing. This antibody is specific to   | •   |
| Notable Publications            | isoforms produced by alternative   | splicing. This antibody is specific to Pubmed ID Journal   | NXN.  |
| Notable Publications            |  |  | NXN.<br>Application                                     |
| Notable Publications            | Author   | Pubmed ID Journal  | NXN.<br>Application<br>Longev WB                        |
| Notable Publications            | Author<br>Claudia Urbainsky  | Pubmed ID Journal<br>30584462 Oxid Med Cell  | NXN.<br>Application<br>Longev WB<br>Med WB              |
| Notable Publications<br>Storage | Author<br>Claudia Urbainsky<br>Kendra M Prutton  | Pubmed ID     Journal       30584462     Oxid Med Cell       35537597     Free Radic Biol       35453370     Antioxidants (Biol) | NXN.<br>Application<br>Longev WB<br>Med WB              |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16128-1-AP (NXN antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using NXN antibody (16128-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594phalloidin (red).



IP result of anti-NXN (IP:16128-1-AP, 4ug; Detection:16128-1-AP 1:500) with HeLa cells lysate 3200ug.



Various lysates were subjected to SDS PAGE followed by western blot with 16128-1-AP (NXN antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.