For Research Use Only

ALG14 Polyclonal antibody

Catalog Number: 16207-1-AP



Purification Method:

IHC 1:50-1:500 IF/ICC 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 16207-1-AP BC011706 GeneID (NCBI): Concentration: 400 ug/ml 199857 **UNIPROT ID:** Source: Rabbit Q96F25 Full Name: Isotype:

asparagine-linked glycosylation 14

Positive Controls:

IF/ICC: A431 cells,

IHC: mouse kidney tissue,

homolog (S. cerevisiae) Immunogen Catalog Number:

AG9178 Calculated MW: 216 aa, 24 kDa

Applications

Tested Applications: IHC, IF/ICC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

ALG14 (UDP-N-Acetylglucosaminyltransferase Subunit ALG14 Homolog) is a protein involved in the process of Nlinked glycosylation, which is essential for protein folding and stability. ALG14 forms a heterodimeric UDP-Nacetylglucosamine transferase (GnTase) with ALG13, which catalyzes a key step in the synthesis of the lipid-linked oligosaccharide (LLO) precursor during N-linked glycosylation (PMID: 36200043).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


human, mouse

PBS with 0.02% sodium azide and 50% glycerol Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 16207-1-AP (ALG14 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A451 cells using ALG14 antibody (16207-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 16207-1-AP (ALG14 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).