For Research Use Only

VIP Polyclonal antibody

Catalog Number: 16233-1-AP 6 Publications



Basic Information

Catalog Number: GenBank Accession Number: 16233-1-AP BC009794

 Size:
 GeneID (NCBI):

 400 µg/ml
 7432

 Source:
 UNIPROT ID:

 Rabbit
 P01282

Isotype: Full Name:

IgG vasoactive intestinal peptide

Immunogen Catalog Number: Calculated MW: AG8933 169 aa, 19 kDa

Applications

Tested Applications:

IHC, IF/ICC, IF-P, ELISA
Cited Applications:
WB. IHC. IF

Species Specificity: human, mouse, rat Cited Species:

mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human colon tissue, rat small intestine tissue, mouse brain tissue, human small intestine tissue

Purification Method:

IHC 1:400-1:1600

IF/ICC 1:50-1:500

IF-P 1:50-1:500

Antigen affinity purification Recommended Dilutions:

IF-P: mouse colon tissue,
IF/ICC: COLO 320 cells,

Background Information

Vasoactive intestinal peptide (VIP), a short peptide containing 28 amino acids belonging to the secretin-glucagon family, is initially isolated from the gastrointestinal tract as a potent vasodilator peptide. VIP was initially identified in normal nervous tissue and neurons and was subsequently recognized as a neurotransmitter widely distributed in various tissues. The wide distribution of VIP determines its involvement in a range of biological activities, such as gut motility, hormonal regulation, circadian rhythms, immune responses, and carcinogenesis. The general physiologic effects of VIP include vasodilation, anti-inflammatory actions, cell proliferation, hormonal secretion, regulation of gastric motility, and smooth muscle relaxation; therefore, VIP has emerged as a promising drug candidate for the treatment of several diseases.

Notable Publications

Author	Pubmed ID	Journal	Application
Muhua Huang	34493162	Pharm Biol	IF
Diane Aguilar	39414787	Nat Commun	
Fatima Aidil-Carvalho	38656542	J Neurosci Res	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16233-1-AP (VIP antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16233-1-AP (VIP antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed COLO 320 cells using 16233-1-AP (VIP antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using VIP antibody (16233-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), Calretinin antibody (66496-1-Ig, Clone: 2D7A9, red).



Immunofluorescent analysis of (4% PFA) fixed mouse colon tissue using VIP antibody (16233-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), Calretinin antibody (66496-1-Ig, Clone: 2D7A9, red).