

For Research Use Only

SFRS5 Polyclonal antibody

Catalog Number: 16237-1-AP **5 Publications**



Basic Information

Catalog Number: 16237-1-AP	GenBank Accession Number: BC018823	Purification Method: Antigen affinity purification
Size: 400 µg/ml	GeneID (NCBI): 6430	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q13243	
Isotype: IgG	Full Name: splicing factor, arginine/serine-rich 5	
Immunogen Catalog Number: AG8988	Calculated MW: 272 aa, 31 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse testis tissue, A549 cells
Cited Applications: WB, IF	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Background Information

SR proteins own a characteristic C-terminal Ser/Arg-rich repeat (RS domain) of variable length and constitute a family of highly conserved nuclear phosphoproteins that can function as both essential and alternative pre-mRNA splicing factors [PMID:7556075]. The binding of SR proteins to pre-mRNA appears to be one of the earliest steps in spliceosome assembly, enhancing the binding of U1 snRNP to the 5' splice site to form a commitment complex [PMID:1577277]. Alternative splicing has a major role in the regulation of gene expression. SFRS5 belongs to the serine/arginine (SR) protein family of regulators of alternative pre-mRNA splicing [PMID:9244433].

Notable Publications

Author	Pubmed ID	Journal	Application
Mi Kyung Park	34879239	Cell Metab	WB
Li-Mei Liang	33905374	JCI Insight	WB
Xuan Gu	37838946	Cell Rep	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 16237-1-AP (SF5S5 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.