For Research Use Only

EXO1 Polyclonal antibody

Catalog Number: 16253-1-AP

Featured Product

9 Publications

BC007491

9156

GeneID (NCBI):

UNIPROT ID:

Q9UQ84 Full Name:

94 kDa

GenBank Accession Number:



Basic Information

Catalog Number: 16253-1-AP

Size:

850 μg/ml Source: Rabbit

Isotype:

Immunogen Catalog Number:

AG9253

Observed MW: 115-120 kDa

exonuclease 1 Calculated MW: **Purification Method:**

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications:

IP, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse **Cited Species:** human, rat

Positive Controls:

WB: HEK-293T cells, Jurkat cells, HeLa cells, Raji cells

IP: mouse thymus tissue,

Background Information

EXO 1, also named as HEX1, is a 846 amino acid protein, which belongs to the XPG/RAD2 endonuclease family. EXO 1 subfamily. EXO1 is highly expressed in bone marrow, testis and thymus and at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach. EXO1 as 5'->3' double-stranded DNA exonuclease, which may also possess a cryptic 3'->5' double-stranded DNA exonuclease activity. EXO1 functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. The calculated molecular weight of EXO1 is $94\,kDa$, but the modified EXO1 protein is about $115\,kDa$.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------------|-----------|-------------------------|-------------|
| Margaret Tammaro | 26420828 | Nucleic Acids Res | WB |
| Christopher B Nelson | 33841503 | Front Genet | WB |
| Xinliang Zou | 35900670 | J Cardiovasc Transl Res | WB |

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16253-1-AP (EXO1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-EXO1 (IP:16253-1-AP, 4ug; Detection:16253-1-AP 1:500) with mouse thymus tissue lysate 2800ug.