For Research Use Only

EXO1/Exonuclease 1 Polyclonal antibody

Catalog Number: 16253-1-AP

Featured Product

15 Publications



Basic Information

Catalog Number: 16253-1-AP Source:

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG9253

GenBank Accession Number:

BC007491 GeneID (NCBI): 9156

UNIPROT ID: Q9UQ84

Full Name: exonuclease 1

Calculated MW: 94 kDa Observed MW: 115-120 kDa Purification Method:

Antigen affinity purification

Recommended Dilutions: WB: 1:2000-1:16000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications: WB, IHC, ChIP Species Specificity:

human, mouse
Cited Species:
human, mouse, rat

Positive Controls:

WB: HEK-293T cells, Jurkat cells, HeLa cells, Raji cells

IP: HEK-293T cells,

Background Information

EXO1, also named as HEX1, is a 846 amino acid protein, which belongs to the XPG/RAD2 endonuclease family. EXO1 subfamily. EXO1 is highly expressed in bone marrow, testis and thymus and at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach. EXO1 as 5'->3' double-stranded DNA exonuclease, which may also possess a cryptic 3'->5' double-stranded DNA exonuclease activity. EXO1 functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. The calculated molecular weight of EXO1 is 94 kDa, but the modified EXO1 protein is about 115 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Margaret Tammaro	26420828	Nucleic Acids Res	WB
Wenbo Liu	36355419	eLife	
Christopher B Nelson	33841503	Front Genet	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16253-1-AP (EXO 1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-EXO1/Exonuclease 1 (IP:16253-1-AP, 4ug; Detection:16253-1-AP 1:2000) with HEK-293T cells lysate 2080 ug.