

PQBP1 Polyclonal antibody

 Catalog Number: 16264-1-AP 3 Publications

Basic Information

Catalog Number: 16264-1-AP Size: 220 µg/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG9329	GenBank Accession Number: BC012358 GeneID (NCBI): 10084 UNIPROT ID: O60828 Full Name: polyglutamine binding protein 1 Calculated MW: 265 aa, 30 kDa Observed MW: 30-32 kDa	Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate IHC 1:500-1:2000 IF 1:50-1:500
---	---	--

Applications

Tested Applications: IF/ICC, IHC, IP, WB, ELISA Cited Applications: WB, IP, IF, IHC Species Specificity: human, mouse, rat Cited Species: human Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	Positive Controls: WB: HeLa cells, mouse brain tissue, mouse skeletal muscle tissue, rat brain tissue IP: mouse brain tissue, IHC: human breast cancer tissue, human colon tissue, rat colon tissue IF: HeLa cells,
--	--

Background Information

PQBP1 (Polyglutamine-binding protein 1) is also named as NPW38 and JM26. PQBP1 senses extrinsic tau 3R/4R proteins by direct interaction and triggers an innate immune response by activating a cyclic GMP-AMP synthase (cGAS)-Stimulator of interferon genes (STING) pathway (PMID: 34782623). PQBP1 is abundantly expressed in the central nervous system during development. PQBP1 is a component of the precatalytic spliceosome (B complex) that is involved in splicing. PQBP1 was predominantly localized in the cell nucleus. PQBP1 is overexpressed in various cancer types including breast cancer, colon cancer, and glioblastoma (PMID: 38342602).


Notable Publications

Author	Pubmed ID	Journal	Application
Hongming Ma	32156810	mBio	WB
Yuqian Shen	33662272	Mol Cell	WB, IP, IF
Neumann Manuela M	22842875	Acta Neuropathol	IHC


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 16264-1-AP (PQBP1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-PQBP1 (IP:16264-1-AP, 4ug; Detection:16264-1-AP 1:500) with mouse brain tissue lysate 3200 ug.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 16264-1-AP (PQBP1 antibody) at dilution of 1:1000 (under 40x lens).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 16264-1-AP (PQBP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).