

CD157 Polyclonal antibody

Catalog Number: 16337-1-AP

Basic Information

Catalog Number: 16337-1-AP	GenBank Accession Number: BC012095	Purification Method: Antigen affinity purification
Size: 200 µg/ml	GeneID (NCBI): 683	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q10588	
Isotype: IgG	Full Name: bone marrow stromal cell antigen 1	
Immunogen Catalog Number: AG9491	Calculated MW: 318 aa, 36 kDa	
	Observed MW: 39 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Species Specificity: human, mouse	WB: A549 cells, IHC: human tonsillitis tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0


Background Information

BST-1 (bone marrow stromal cell antigen 1), also named as CD157 or ADP-ribosyl cyclase 2, is a glycosyl phosphatidylinositol-anchored protein that stimulates pre-B-cell growth and has a adenosine diphosphate (ADP)-ribosyl cyclase and cyclic ADP-ribose (cADPR) hydrolase activity (PMID: 9473467, 11866528). The two enzymatic activities are responsible for the synthesis and hydrolysis of cADPR, a novel second messenger of calcium release from intracellular calcium stores. BST-1 has a calculated molecular mass of 30-36 kDa and an apparent molecular mass of 40-46 kDa due to the glycosylation (PMID: 8941363, 10491089).


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 16337-1-AP (CD157 antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 16337-1-AP (CD157 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 16337-1-AP (CD157 antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).