

For Research Use Only

TXNRD2 Polyclonal antibody

Catalog Number: 16360-1-AP

13 Publications



Basic Information

Catalog Number:

16360-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG8367

GenBank Accession Number:

BC007489

GeneID (NCBI):

10587

UNIPROT ID:

Q9NNW7

Full Name:

thioredoxin reductase 2

Calculated MW:

524 aa, 56 kDa

Observed MW:

54 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IHC 1:50-1:500

Applications

Tested Applications:

IHC, WB, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, rat liver tissue, MCF-7 cells, HeLa cells

IHC : human liver tissue,

Background Information

Human mitochondrial thioredoxin reductase (TXNRD2) is a selenocysteine-containing enzyme essential for mitochondrial oxygen radical scavenging. Three mammalian thioredoxin reductases exist; a cytosolic (TXNRD1), a mitochondrial (TXNRD2), and a testis-specific thioredoxin reductase (TXNRD3). (PMID: 21247928, PMID: 26199228)

Notable Publications

Author	Pubmed ID	Journal	Application
B Chellan	33122797	Sci Rep	WB
Caitlin E. Miller	36358489	Antioxidants (Basel)	WB
Tai-Jung Lee	36336122	Arch Biochem Biophys	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16360-1-AP (TXNRD2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16360-1-AP (TXNRD2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).