## For Research Use Only

## **UBQLNL** Polyclonal antibody

Catalog Number:16400-1-AP 1 Publications



**Basic Information** 

Catalog Number: 16400-1-AP GenBank Accession Number: BC012183

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

900 ug/ml Source: 143630 UNIPROT ID: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

Rabbit Isotype:

Q8IYU4 Full Name: ubiquilin-like protein lysate

IgG

Calculated MW:

IHC 1:50-1:500 IF/ICC 1:10-1:100

Immunogen Catalog Number: AG9496

475 aa, 53 kDa Observed MW:

70 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IP, ELISA

Positive Controls:

WB: human brain tissue, Jurkat cells

Cited Applications:

IP: Jurkat cells.

WB, IF

IHC

Species Specificity: human, mouse IHC: human colon tissue,

IF/ICC: A431 cells,

Cited Species:

mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Background Information** 

UBQLNL also named as Ubiquilin like protein, encodes a 475 amino acid protein. The specific function of it is still non-known. UBQLNL may have an association with type 2 diabetes, but the mechanism is supposed be studied in the further research. The subcellular localization of UBQLNL is nucleus. The expression of it is detected in some tissues, such as brain ,testis, heart and liver.

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Li Luo	31059032	Mol Med Rep	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




human brain tissue were subjected to SDS PAGE followed by western blot with 16400-1-AP (UBQLNL antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of A431 cells using 16400-1-AP (UBQLNL antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16400-1-AP (UBQLNL antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using UBQLNL antibody (16400-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



IP result of anti-UBQLNL (IP:16400-1-AP, 4ug; Detection:16400-1-AP 1:500) with Jurkat cells lysate 3200ug.