For Research Use Only

Aquaporin 4 Polyclonal antibody, PBS Only



Purification Method:

Antigen affinity purification

Catalog Number:16473-1-PBS

Featured Product

Basic Information

Catalog Number:

16473-1-PBS BC022286

Source: GeneID (NCBI):

Rabbit 36

 Isotype:
 UNIPROT ID:

 IgG
 P55087

 Immunogen Catalog Number:
 Full Name:

Calculated MW: 323 aa, 35 kDa Observed MW: 35-37 kDa, 32-34 kDa

Aquaporin 4

GenBank Accession Number:

Applications

Tested Applications:

AG9561

WB, IHC, IF-P, IF-Fro, IP, ELISA

Species Specificity: human, mouse, rat

Background Information

Aquaporins are specialized water transport channels in plasma membranes of water-permeable tissues. Aquaporin-4 (AQP4) is the most abundant water channel in the human central nervous system and is important to fluid movements in brain. Aquaporin-4 exists as two isoforms, a long (M1) isoform with translation initiation at Met-1, and a shorter (M23) isoform with translation initiation at Met-23, with molecular weights around 35-37 kDa and 32-34 kDa, respectively.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only, pH7.3


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16473-1-AP (Aquaporin 4 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16473-1-AP (Aquaporin 4 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16473-1-AP (Aquaporin 4 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 16473-1-AP (Aquaporin 4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.




IP result of anti-Aquaporin 4 (IP:16473-1-AP, 4ug; Detection:16473-1-AP 1:300) with mouse heart tissue lysate 4000ug. This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Aquaporin 4 antibody (16473-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L), GFAP antibody (60190-1-Ig, Clone: 4BZE10, red). This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using Aquaporin 4 antibody (16473-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), GFAP antibody (60190-1-Ig, Clone: 4B2E10, red). This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using Aquaporin 4 antibody (16473-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 16473-1-PBS in a different storage buffer formulation