For Research Use Only

AXIN1 Polyclonal antibody Catalog Number: 16541-1-AP Featured Product

Featured Product





Basic Information	Catalog Number: 16541-1-AP	GenBank Accession Number: BC044648	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	600 ug/ml	8312	WB 1:1000-1:6000
	Source: Rabbit	UNIPROT ID: 015169	IHC 1:50-1:500
	Isotype: IgG	Full Name: axin 1	
	Immunogen Catalog Number: AG9858	Calculated MW: 826aa,92 kDa; 862aa,95 kDa	
		Observed MW: 100-110 kDa	
Applications	Tested Applications: WB, IHC, ELISA	Positive Controls:	
	Cited Applications: WB, IHC, IP	WB : HeLa cells, HT-1080 cells, HEK-293T cells IHC : human colon tissue,	
	Species Specificity: human		
	Cited Species: human, rat, chicken		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Informatior	negatively regulate Wnt signalin HIPK2-TP53 complex forms. The c undergoes poly(ADP-ribosy)lation leads to its degradation and subso nuclear accumulation during Wnt 295aa, which shows at around 40- the Wnt signaling pathway to pro	g. AXIN1 is likely to function as a tum complex also controls cell growth, apo n by tankyrase TNKS and TNKS2 follow equent activation of Wnt signaling. It signaling. Recent researches find tha -55 kDa by Western Blot. AXIN1-295aa mote GC tumorigenesis and progressi	ptosis and development. Like AXIN2, AXIN wed by unbiquitination by RNF146 which s deubiquitination by USP34 is important for t CircAXIN1 encodes a novel protein, AXIN1 a functions as an oncogenic protein, activat
	negatively regulate Wnt signalin HIPK2-TP53 complex forms. The c undergoes poly(ADP-ribosy)lation leads to its degradation and subso nuclear accumulation during Wnt 295aa, which shows at around 40- the Wnt signaling pathway to pro for GC. Proteintech's AXIN1 antibo	g. AXIN1 is likely to function as a tum complex also controls cell growth, apo n by tankyrase TNKS and TNKS2 follow equent activation of Wnt signaling. It signaling. Recent researches find tha -55 kDa by Western Blot. AXIN1-295aa mote GC tumorigenesis and progressi	or suppressor. Under UV irradiation, AXIN1- potosis and development. Like AXIN2, AXIN wed by unbiquitination by RNF146 which s deubiquitination by USP34 is important for t CircAXIN1 encodes a novel protein, AXIN1 a functions as an oncogenic protein, activat ion, suggesting a potential therapeutic targ antibdy raised against the N-terminus of
	negatively regulate Wnt signalin HIPK2-TP53 complex forms. The c undergoes poly(ADP-ribosy)lation leads to its degradation and subso nuclear accumulation during Wnt 295aa, which shows at around 40- the Wnt signaling pathway to pro for GC. Proteintech's AXIN1 antibo human AXIN1.	g. AXIN1 is likely to function as a tum complex also controls cell growth, apo n by tankyrase TNKS and TNKS2 follow equent activation of Wnt signaling. It signaling. Recent researches find tha -55 kDa by Western Blot. AXIN1-295aa mote GC tumorigenesis and progressi ody 16541-1-AP is a rabbit polyclonal Pubmed ID Journal	or suppressor. Under UV irradiation, AXIN1- potosis and development. Like AXIN2, AXIN wed by unbiquitination by RNF 146 which a deubiquitination by USP34 is important for t CircAXIN1 encodes a novel protein, AXIN1 a functions as an oncogenic protein, activat ion, suggesting a potential therapeutic targ antibdy raised against the N-terminus of Application
	negatively regulate Wnt signalin HIPK2-TP53 complex forms. The c undergoes poly(ADP-ribosy)lation leads to its degradation and subse nuclear accumulation during Wnt 295aa, which shows at around 40- the Wnt signaling pathway to pro for GC. Proteintech's AXIN1 antibo human AXIN1.	g. AXIN1 is likely to function as a tum complex also controls cell growth, apo n by tankyrase TNKS and TNKS2 follow equent activation of Wnt signaling. It signaling. Recent researches find tha -55 kDa by Western Blot. AXIN1-295aa mote GC tumorigenesis and progressi ody 16541-1-AP is a rabbit polyclonal Pubmed ID Journal	or suppressor. Under UV irradiation, AXIN1 optosis and development. Like AXIN2, AXIN wed by unbiquitination by RNF 146 which a deubiquitination by USP34 is important for t CircAXIN1 encodes a novel protein, AXIN1 a functions as an oncogenic protein, activat ion, suggesting a potential therapeutic targ antibdy raised against the N-terminus of Application
Background Information	negatively regulate Wnt signalin HIPK2-TP53 complex forms. The c undergoes poly(ADP-ribosy)lation leads to its degradation and subsr nuclear accumulation during Wnt 295aa, which shows at around 40- the Wnt signaling pathway to pro for GC. Proteintech's AXIN1 antibo human AXIN1.	g. AXIN1 is likely to function as a tum complex also controls cell growth, apo n by tankyrase TNKS and TNKS2 follox equent activation of Wnt signaling. It signaling. Recent researches find that -55 kDa by Western Blot. AXIN1-295aa mote GC tumorigenesis and progressi ody 16541-1-AP is a rabbit polyclonal Pubmed ID Journal 36305172 Environ Toxicol	or suppressor. Under UV irradiation, AXIN1 optosis and development. Like AXIN2, AXIN wed by unbiquitination by RNF146 which s deubiquitination by USP34 is important for t CircAXIN1 encodes a novel protein, AXIN1 a functions as an oncogenic protein, activat ion, suggesting a potential therapeutic targ antibdy raised against the N-terminus of Application WB WB


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16541-1-AP (AXIN1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16541-1-AP (AXIN1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 16541-1-AP (AXIN1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.