For Research Use Only

SDS Polyclonal antibody

Catalog Number: 16594-1-AP



Basic Information

Catalog Number:

GenBank Accession Number:

16594-1-AP

BC020750

Size:

GeneID (NCBI):

700 µg/ml

10993

Source:

UNIPROT ID:

Rabbit

P20132

Isotype:

Full Name:

IgG serine dehydratase
Immunogen Catalog Number: Calculated MW:
AG9704 218 aa, 23 kDa
Observed MW:
35 kDa; 70 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500

Applications

Tested Applications: IHC, WB, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse liver tissue, IHC: human liver tissue,

Background Information

L-Serine dehydratase (SDS; SDH) catalyzes the pyridoxal phosphate (PLP) dependent deamination of l-serine to yield pyruvate. In mammals, SDH is found predominantly in the liver. Extensive studies have been carried out on SDH from rat and the enzyme has been found to play an important role in gluconeogenesis; its activity is induced by the consumption of high-protein diets, starvation and other treatments. The enzyme exists as a dimer of identical subunits, with each subunit exhibiting a bilobal architecture. (PMID: 25380533, PMID: 21878319, PMID: 14646100)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16594-1-AP (SDS antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 16594-1-AP (SDS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 16594-1-AP (SDS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).