For Research Use Only

LIPH Polyclonal antibody

Catalog Number: 16602-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number:

16602-1-AP

BC064941

Concentration:

500 ug/ml

Source:

Rabbit

UNIPROT ID:

Rabbit

Q8WWY8

Isotype:

IgG

GenelD (NCBI):

Full Name:

Igg lipase, member H

Immunogen Catalog Number: Calculated MW: AG9775 51 kDa

Observed MW: 50-55 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000

IF/ICC 1:10-1:100

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, ELISA Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

IF/ICC: A549 cells,

WB: A549 cells, COLO 320 cells, mouse lung tissue

IP: mouse lung tissue,
IHC: human pancreas tissue,

Background Information

LIPH(Lipase member H) is also named as LPDLR, MPAPLA1, PLA1B. It is a phosphatidic acid-selective phospholipase A1 (PLA1) that produces 2-acyl lysophosphatidic acid (LPA) and belongs to the AB hydrolase superfamily and Lipase family.LIPH has a 12-residue lid region, which likely covers a catalytic pocket, and 4 potential N-linked glycosylation sites. Defects in LIPH are the cause of hypotrichosis type 7 (HYPT7).

Notable Publications

Author	Pubmed ID	Journal	Application
Hisako Ishimine	26341494	Tumour Biol	IHC
Yixiao Zhang	32618099	J Cell Mol Med	WB
Yasuhiro Seki	24380866	Biochem Biophys Res Commun	ELISA,WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




A549 cells were subjected to SDS PAGE followed by western blot with 16602-1-AP (LIPH antibody) at dilution of 1:500 incubated at room temperature for 15 hours.




Immunohistochemical analysis of paraffinembedded human pancreas using 16602-1-AP (LIPH antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human pancreas using 16602-1-AP (LIPH antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-LIPH (IP:16602-1-AP, 4ug; Detection:16602-1-AP 1:500) with mouse lung tissue lysate 4000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed A549 cells using 16602-1-AP (LIPH antibody) at dilution of 1:25 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).