## For Research Use Only

# OTUD7B Polyclonal antibody

Catalog Number: 16605-1-AP

Featured Product

24 Publications



**Basic Information** 

Catalog Number:

16605-1-AP

BC020622

Concentration:

550 µg/ml

56957

Source:

Rabbit

Q6GQQ9

Isotype:

GenBank Accession Number:

BC020622

GeneID (NCBI):

56957

UNIPROT ID:

Q6GQQ9

Full Name:

gG OTU domain containing 7B

Immunogen Catalog Number: Calculated MW:

AG9788 427aa,48 kDa; 843aa,93 kDa

Observed MW: 100 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

# **Applications**

Tested Applications:
WB, IP, IHC, ELISA
Cited Applications:
WB, IHC, IF, IP, CoIP
Species Specificity:
human, mouse, rat
Cited Species:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: Jurkat cells, IP: Jurkat cells,

IHC: human kidney tissue, human heart tissue, mouse kidney tissue, mouse heart tissue

# **Background Information**

OUTD7B is a member of the A20 family of the deubiquitinating cysteine proteases, which has deubiquitinating activity that is directed towards 'Lys-48' or 'Lys-63'-linked polyubiquitin chains. It plays a key role in the negative feedback regulation of NF-  $\kappa$  B signaling in response to multiple stimuli and has been described as central gatekeeper in inflammation and immunity. The calculated molecular weight of OTUD7B is 93 kDa, but the modified OTUD7B is about 100 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Dan-Dan Lin	31572671	Front Oncol	IHC
Shuang Feng	28882872	Arterioscler Thromb Vasc Biol	WB,IF
Chun-Ping Cui	30405104	Nat Commun	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 16605-1-AP (OTUD7B antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney using 16605-1-AP (OTUD7B antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-OTUD7B (IP:16605-1-AP, 4ug; Detection:16605-1-AP 1:800) with Jurkat cells lysate 2500ug.