

For Research Use Only

# AMZ2 Polyclonal antibody

Catalog Number: 16664-1-AP



## Basic Information

Catalog Number:

16664-1-AP

Size:

700 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10141

GenBank Accession Number:

BC056271

GeneID (NCBI):

51321

UNIPROT ID:

Q86W34

Full Name:

archaelysin family metallopeptidase

2

Calculated MW:

360 aa, 41 kDa

Observed MW:

41 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF/ICC 1:50-1:500

## Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB : HepG2 cells, human brain tissue

IHC : human liver tissue, human heart tissue, human placenta tissue, human testis tissue, human skin tissue, human brain tissue, human spleen tissue, human ovary tissue

IF/ICC : HepG2 cells,

## Background Information

AMZ2(archaelysin family metallopeptidase 2) is a zinc metalloprotease. Its calculated molecular weight and observed molecular weight are both 41kDa.

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16664-1-AP (AMZ2 antibody) at dilution of 1:100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 16664-1-AP (AMZ2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1x10<sup>6</sup> HepG2 cells were intracellularly stained with 0.4 ug AMZ2 Polyclonal antibody (16664-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



HepG2 cells were subjected to SDS PAGE followed by western blot with 16664-1-AP (AMZ2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human testis using 16664-1-AP (AMZ2 antibody) at dilution of 1:50 (under 10x lens).