For Research Use Only

CD3 Delta Polyclonal antibody

Catalog Number: 16669-1-AP 9 Publications



Basic Information

Catalog Number: 16669-1-AP

Size: 600 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG10148

171 aa, 19 kDa Observed MW:

GenBank Accession Number:

BC070321 GeneID (NCBI):

UNIPROT ID: P04234 Full Name:

CD3d molecule, delta (CD3-TCR complex)

Calculated MW:

20-25 kDa

Applications

Tested Applications: WB, IF-P, IP, ELISA **Cited Applications:** WB, IHC, IF, IP

Species Specificity: human, mouse, rat Cited Species:

human, mouse

Positive Controls:

WB: Jurkat cells, mouse thymus tissue, rat thymus

Purification Method:

WB 1:1000-1:4000

protein lysate

IF-P 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

tissue, MOLT-4 cells IP: Jurkat cells,

IF-P: human tonsillitis tissue,

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T $lymphocytes\ consists\ of\ either\ a\ TCR\ alpha/beta\ or\ TCR\ gamma/delta\ heterodimer\ coexpressed\ at\ the\ cell\ surface$ with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. CD3 delta/CD3D is a single-pass type I membrane protein which consists of an extracellular domain of 84 amino acids, a transmembrane region, and a cytoplasmic domain of 45 amino acids. Defects in CD3 delta are a cause of severe combined immunodeficiency.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuji Fukushima	36130493	Cell Rep	WB,IP
Marie Klintman	27179111	Clin Cancer Res	
Binghao Li	29735547	Cancer Res	IHC

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16669-1-AP (CD3 Delta antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 Delta antibody (16669-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 Delta antibody (16669-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-CD3 Delta (IP:16669-1-AP, 4ug; Detection:16669-1-AP 1:800) with Jurkat cells lysate 2800ug.