## For Research Use Only

## CKAP4 Polyclonal antibody

Catalog Number: 16686-1-AP 52 Publications



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 16686-1-AP
 BC082972

 Source:
 GeneID (NCBI):

 Rabbit
 10970

 Isotype:
 UNIPROT ID:

 IgG
 Q07065

 Immunogen Catalog Number:
 Full Name:

 AG10019
 cytoskeleton-associated protein 4

Calculated MW: 602 aa, 66 kDa Observed MW: 63 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:50-1:500 IF/ICC: 1:50-1:500

WB: 1:5000-1:50000

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IF, IP, CoIP
Species Specificity:
human, mouse, rat
Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: A431 cells, rat kidney tissue, HeLa cells, HepG2

cells, NIH/3T3 cells
IP: HeLa cells,

IHC: human kidney tissue, human liver tissue

IF/ICC: HepG2 cells,

**Background Information** 

CKAP4, also named as Climp-63, is a 63 kDa cytoskeleton-linking membrane protein. It is the partner of triadin. CKAP4 is responsible for this association of triads and microtubules.

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Gui-Feng Lu	30288057	Onco Targets Ther	WB,IHC
Svenja Zielke	33111629	Autophagy	WB,IF
Michael L van de Weijer	33065021	Mol Cell	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16686-1-AP (CKAP4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 16686-1-AP (CKAP4 antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.



IP result of anti-CKAP4 (IP:16686-1-AP, 4ug; Detection:16686-1-AP 1:1000) with HeLa cells lysate 2000ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CKAP4 antibody (16686-1-AP) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).