## For Research Use Only

## CKAP4 Polyclonal antibody

Catalog Number: 16686-1-AP

38 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

BC082972

Size:

GeneID (NCBI):

400 µ g/ml

10970

Source:

UNIPROT ID:

Rabbit

Q07065

Isotype:

GenBank Accession Number:

BC082972

UNCBI):

Q07065

cytoskeleton-associated protein 4

Immunogen Catalog Number:Calculated MW:AG10019602 aa, 66 kDaObserved MW:

63 kDa

**Applications** 

Tested Applications: IF/ICC, IHC, IP, WB, ELISA

Cited Applications: WB, IP, IF, IHC, CoIP Species Specificity: human, mouse, rat

Cited Species: human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A431 cells, rat kidney tissue, HeLa cells, HepG2

**Purification Method:** 

WB 1:5000-1:50000

protein lysate

IHC 1:50-1:500

IF 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

cells, NIH/3T3 cells
IP: HeLa cells,

IHC: human kidney tissue, human liver tissue

IF: HepG2 cells,

**Background Information** 

CKAP4, also named as Climp-63, is a 63 kDa cytoskeleton-linking membrane protein. It is the partner of triadin. CKAP4 is responsible for this association of triads and microtubules.

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Gui-Feng Lu	30288057	Onco Targets Ther	WB,IHC
Svenja Zielke	33111629	Autophagy	WB,IF
Michael L van de Weijer	33065021	Mol Cell	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16686-1-AP (CKAP4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 16686-1-AP (CKAP4 antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CKAP4 antibody (16686-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-CKAP4 (IP:16686-1-AP, 4ug; Detection:16686-1-AP 1:1000) with HeLa cells lysate 2000ug.