

For Research Use Only

NKD2 Polyclonal antibody, PBS Only

Catalog Number: 16699-1-PBS

Featured Product



Basic Information

Catalog Number:

16699-1-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10094

GenBank Accession Number:

BC012176

GeneID (NCBI):

85409

UNIPROT ID:

Q969F2

Full Name:

naked cuticle homolog 2 (Drosophila)

Calculated MW:

50 kDa

Observed MW:

59 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

Naked2 (protein naked cuticle homolog 2), encoded by NKD2 gene, was identified as a cell autonomous antagonist of the canonical Wnt signaling pathway. Naked2 acts as a cargo recognition and targeting protein to ensure proper delivery and fusion of TGF- α -containing vesicles to a distinct region at the basolateral surface of polarized epithelial cells (PMID: 17553928). EGFR-independent action of TGF- α protects Naked2 from proteasomal degradation, thus ensuring its delivery to the basolateral surface (PMID: 18757723). Myristoylated Naked2 antagonizes Wnt-beta-catenin activity possibly by degrading Dishevelled-1 at the plasma membrane (PMID: 20177058).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 16699-1-AP (NKD2 antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 16699-1-PBS in a different storage buffer formulation.



WB result of NKD2 antibody (16699-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-NKD2 transfected HEK-293 cells. This data was developed using the same antibody clone with 16699-1-PBS in a different storage buffer formulation.