For Research Use Only

N4BP3 Polyclonal antibody

Catalog Number: 16733-1-AP

Featured Product

3 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 16733-1-AP
 BC053323

 Source:
 GeneID (NCBI):

 Rabbit
 23138

 Isotype:
 UNIPROT ID:

IgG 015049
Immunogen Catalog Number: Full Name:

AG10194 Nedd4 binding protein 3

Calculated MW: 544 aa, 60 kDa Observed MW: 60-65 kDa Purification Method:

Antigen affinity purification Recommended Dilutions:

WB: 1:1000-1:6000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:50-1:500 IF/ICC: 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC, IP Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse brain tissue, rat brain tissue

IP: mouse brain tissue,

IHC: mouse kidney tissue, human cervical cancer tissue, human kidney tissue, mouse brain tissue

IF/ICC: HEK-293 cells,

Background Information

N4BP3, also named as KIAAO341, a Fezzin family member, during nerve cell development. It is crucial for axonal and dendritic branching in developing neurons. It binds the protein NEDD4 and interacts with 14-3-3 proteins. N4BP3 is present in early neurites, growth cones, axons and dendrites of developing primary hippocampal neurons.

Notable Publications

Author	Pubmed ID	Journal	Application
Chen Wang	34880843	Front Microbiol	WB,IP
Wang Jiang	40487287	J Inflamm Res	WB,IP
Wang Jiang	39420190	Cell Death Discov	WB,IP,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16733-1-AP (N4BP3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 16733-1-AP (N4BP3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 16733-1-AP (N4BP3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using N4BP3 antibody (16733-1-AP) at dilution of 1:200 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



IP result of anti-N4BP3 (IP:16733-1-AP, 4ug; Detection:16733-1-AP 1:2000) with mouse brain tissue lysate 1800 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16733-1-AP (N4BP3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using N4BP3 antibody (16733-1-AP) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).