For Research Use Only

PCK1/2 Polyclonal antibody

Catalog Number: 16754-1-AP

Featured Product 78 Publications



Purification Method:

WB 1:5000-1:50000

protein lysate

IF/ICC 1:50-1:500

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 16754-1-AP BC023978 Concentration: GeneID (NCBI): 650 ug/ml 5105 **UNIPROT ID:** Source: Rabbit P35558 Full Name: Isotype:

> phosphoenolpyruvate carboxykinase 1 (soluble)

Calculated MW: 622 aa, 69 kDa Observed MW: 66-69 kDa

Applications

Tested Applications: WB, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP Species Specificity: human, mouse, rat

Cited Species:

Immunogen Catalog Number:

AG10261

human, mouse, rat, pig, bovine, goat

Positive Controls:

WB: mouse liver tissue, rat liver tissue

IP: HepG2 cells, IF/ICC: HepG2 cells,

Background Information

PCK1(Phosphoenolpyruvate carboxykinase, cytosolic) is also named as PEPCK1 and belongs to the $phosphoenol pyruvate\ carboxykin as e\ [GTP]\ family.\ It\ catalyzes\ the\ formation\ of\ phosphoenol pyruvate\ from$ oxaloacetate, with the release of carbon dioxide and GDP. It is also a main control point for the regulation of gluconeogenesis. In eukaryotes there are two isozymes: a cytoplasmic one and a mitochondrial one. Defects in PCK1 are the cause of cytosolic phosphoenolpyruvate carboxykinase deficiency (C-PEPCKD). This antibody has weak cross-reactivity with PCK2 protein due to high sequence homology between PCK1 and PCK2.

Notable Publications

Author	Pubmed ID	Journal	Application
Yong Zhang	32970960	Can J Physiol Pharmacol	WB
Teresa W-M Fan	36150727	J Immunol	
Wei Xu	32935679	Food Funct	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16754-1-AP (PCK1/2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-PCK1/2 (IP:16754-1-AP, 4ug; Detection:16754-1-AP 1:500) with HepG2 cells lysate 3600ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PCK1/2 antibody (16754-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).