For Research Use Only

NEDD8 Polyclonal antibody Catalog Number: 16777-1-AP Featured Product 8

Featured Product 8 Publications





Basic Information	Catalog Number: 16777-1-AP	GenBank Accession Number: BC104664		Purification Method: Antigen affinity purific	ation
	Concentration:	GenelD (NCE	31):	Recommended Dilutions:	
	600 ug/ml	4738		WB 1:500-1:2000 IHC 1:250-1:1000	
	Source: Rabbit	UNIPROT ID: Q15843	UNIPROTID.		HC 1:250-1:1000
	Isotype: IgG Immunogen Catalog Number: AG10154		-		
		Calculated M			
		Observed MV 6-10 kDa	N:		
Applications	Tested Applications: WB, IHC, ELISA		Positive Controls:		
	Cited Applications: WB, IHC, IF		WB : mouse brain tissue, human heart tissue IHC : human prostate cancer tissue,		
	Species Specificity: human, mouse, rat, monkey				
	Cited Species: human, mouse, rat				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	NEDD8 (neural precursor cell-expressed developmentally down-regulated gene 8) is a small ubiquitin-like prote that shares 60% sequence identity and 80% homology with ubiquitin [PMID:9353319]. NEDD8 is conjugated to substrate proteins in a process known as neddylation. NEDD8 forms a thioester bond with APPBP1-Uba3, the E1 enzyme. Activated NEDD8 then transfers to Ubc12, the E2 enzyme. Ultimately, E2 loaded with NEDD8 binds a substrate protein lysine residue and forms an isopeptide bond by E3 ligase [PMID:18802447].				
Notable Publications	Author	Pubmed ID	Journal	A	pplication
	Kaile Zhang	32478052	Front Bioeng Biotechnol		/B
	Lu Boxun B	23525043	Nat Neurosci		
	Gonca Bayraktar	32726795	Neuropsychopharmacol	ogy II	F,WB

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse brain tissue were subjected to SDS PAGE followed by western blot with 16777-1-AP (NEDD8 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 16777-1-AP (NEDD8 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).