

For Research Use Only

DCTD Polyclonal antibody

Catalog Number: 16784-1-AP

1 Publications



Basic Information

Catalog Number:

16784-1-AP

Size:

750 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10216

GenBank Accession Number:

BC001286

GeneID (NCBI):

1635

UNIPROT ID:

P32321

Full Name:

dCMP deaminase

Calculated MW:

178 aa, 20 kDa

Observed MW:

20 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human

Positive Controls:

WB : human brain tissue, PC-3 cells

IF/ICC : HepG2 cells,

Background Information

DCTD(Deoxycytidylate deaminase) is also named as dCMP deaminase and belongs to the cytidine and deoxycytidylate deaminase family. It catalyzes the deamination of dCMP to dUMP, thus providing the nucleotide substrate for thymidylate synthase. Control of deaminase activity at this juncture in deoxyribonucleotide metabolism is determined by the ratio of dCTP to dTTP in the cell, since the enzyme is allosterically activated by dCTP and inhibited by dTTP(PMID:7685356). It has 2 isoforms produced by alternative splicing and can exist as a dimer(PMID:8798492).

Notable Publications

Author	Pubmed ID	Journal	Application
Nao Yoshida-Sakai	34913485	Int J Cancer	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 16784-1-AP (DCTD antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using DCTD antibody (16784-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).