### For Research Use Only

# MPP8 Polyclonal antibody

Catalog Number: 16796-1-AP

**Featured Product** 

28 Publications



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 16796-1-AP
 BC003542

 Size:
 GeneID (NCBI):

 700 μg/ml
 54737

 Source:
 UNIPROT ID:

 Rabbit
 Q99549

Isotype: Full Name:
IgG M-phase phosphoprotein 8

Immunogen Catalog Number: Calculated MW:

AG10268 Refuel AG1026

97-105 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
IF/ICC 1:50-1:500

**Applications** 

Tested Applications: WB, IF/ICC, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP, chIP Species Specificity: human, mouse, rat Cited Species:

human, mouse

#### **Positive Controls:**

WB: HEK-293 cells, HeLa cells, HepG2 cells, MCF-7 cells, U-251 cells, NIH/3T3 cells, mouse brain tissue, rat brain tissue

IP: HEK-293 cells,
IF/ICC: U2OS cells,

# **Background Information**

The M-phase phosphoprotein 8 (MPP8) was firstly identified to coimmunoprecipitate with the RanBPM-comprised large protein complex [PMID:19956676]. It also binds to methylated H3K9 and recruited the H3K9 methyltransferases GLP and ESET, as well as DNA methyltransferase 3A (DNMT3A) to the promoter of the E-cadherin gene, a key regulator of tumor cell growth and epithelial-to-mesenchymal transition (EMT) [PMID:18726070]. This binding event of MPP8-H3K9me modulates the expression of E-cadherin, a key regulator of cell-cell adhesion with implications in embryonic development, tissue morphogenesis and homeostasis, together with H3K9 methyltransferases GLP and ESET and DNA methyltransferase 3a (Dnmt3a) [PMID: 18726070].

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Paola Cruz-Tapias	31557926	Genes (Basel)	WB,IP,chIP
Leonid Yurkovetskiy	30297740	Nat Microbiol	WB
Takeshi Tsusaka	30286792	Epigenetics Chromatin	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16796-1-AP (MPP8 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-MPP8 (IP:16796-1-AP, 4ug; Detection:16796-1-AP 1:20000) with HEK-293 cells lysate 1400 ug.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using MPP8 antibody (16796-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).