For Research Use Only

GFAP Polyclonal antibody

Catalog Number: 16825-1-AP

Featured Product

620 Publications



Basic Information

Catalog Number: GenBank Accession Number: 16825-1-AP BC013596

Concentration: GeneID (NCBI): 2670

Source: UNIPROT ID: Rabbit P14136

Isotype: Full Name:

gG glial fibrillary acidic protein

Immunogen Catalog Number: Calculated MW:
AG10423 432 aa, 50 kDa
Observed MW:

45-50 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IF-Fro, ELISA

Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, rabbit, monkey, zebrafish,

hamster, goat, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Purification Method:**

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:10000 IHC 1:2500-1:10000 IF-P 1:50-1:500 IF-Fro 1:500-1:2000 IF/ICC 1:50-1:500

Positive Controls:

WB: mouse brain tissue, U-251 cells, rat brain tissue

IHC: mouse brain tissue, Alzheimer mouse

IF-P: rat brain tissue, mouse brain tissue

IF-Fro: mouse brain tissue,
IF/ICC: A172 cells,

Background Information

GFAP (Glial fibrillary acidic protein), an intermediate-filament (IF) protein, is specifically expressed in cells of astroglial lineage and is widely used to mark astroglia in the brain. It is also used as a marker for intracranial and intraspinal tumors arising from astrocytes.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|-------------------|-------------|
| Yi-Na Zhang | 36168082 | Transl Stroke Res | IF |
| Yihua He | 34585253 | J Radiat Res | IF |
| Shuisheng Yu | 34646136 | Front Pharmacol | IF |

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded Hippocampus tissue slide from Alzheimer Mouse Model using 16825-1-AP (GFAP antibody) at dilution of 1:50000 (under 10x lens). Data from NeuroScience Associates, Inc.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using 16825-1-AP (GFAP antibody) at dilution of 1:1000 and Coralite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16825-1-AP (GFAP antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16825-1-AP (GFAP antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of GFAP antibody (16825-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GFAP transfected U-251 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 16825-1-AP (GFAP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse brain tissue using GFAP antibody (16825-1-AP) at dilution of 1:200 and Coralite®594-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-4).



Immunofluorescent analysis of (-20°C Ethanol) fixed A 172 cells using GFAP antibody (16825-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).