# For Research Use Only

# AP1B1 Polyclonal antibody

Catalog Number: 16932-1-AP

5 Publications



**Purification Method:** 

WB 1:500-1:2400

IF/ICC 1:50-1:500

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number: 16932-1-AP Size:

750 µg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

AG10417

**Tested Applications:** 

IF/ICC, IP, WB, ELISA Cited Applications:

Species Specificity:

human Cited Species: human

GenBank Accession Number:

BC046242 GeneID (NCBI):

**UNIPROT ID:** Q10567

Full Name:

adaptor-related protein complex 1,

beta 1 subunit Calculated MW: 919 aa, 101 kDa

Observed MW: 101 kDa

Positive Controls:

WB: HEK-293 cells, IP: HEK-293 cells, IF/ICC: HepG2 cells,

# **Background Information**

AP1B1 is a subunit of the adaptor complex AP-1 and is a member of the adaptin protein family. Adaptor protein (AP) complexes are cytosolic heterotetramers that mediate the sorting of membrane proteins in the secretory and endocytic pathways. AP-1 is found at the cytoplasmic face of coated vesicles located at the Golgi complex, where it mediates both the recruitment of clathrin to the membrane and the recognition of sorting signals within the cytosolic tails of transmembrane receptors.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Arunas Kvainickas	28935632	J Cell Biol	WB
Lynn M Boyden	31630788	Am J Hum Genet	WB
Antoine Rebendenne	35879413	Nat Genet	WB

# Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


# Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 16932-1-AP (AP1B1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-AP1B1 (IP:16932-1-AP, 4ug; Detection:16932-1-AP 1:300) with HEK-293 cells lysate 2320 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 16932-1-AP (AP1B1 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).