For Research Use Only

RNF10 Polyclonal antibody

Catalog Number: 16936-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: 16936-1-AP GenBank Accession Number: BC016622

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

650 ug/ml

9921

WB 1:500-1:2000

Source: Rabbit UNIPROT ID: Q8N5U6 IP 0.5-4.0 ug for 1.0-3.0 mg of total

Isotype:

Full Name:

ring finger protein 10

protein lysate IHC 1:250-1:1000 IF/ICC 1:50-1:500

Immunogen Catalog Number: AG10433

Calculated MW: 811 aa, 90 kDa

811 aa, 90 kD

Observed MW: 90 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Α

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species: human, mouse, rat Positive Controls:

WB: PC-3 cells, Jurkat cells, human brain tissue

IP: PC-3 cells,

IHC: mouse brain tissue, human brain tissue

IF/ICC: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

RING finger (C3HC4-type zinc finger) is a variant zinc finger motif present in a new family of proteins including transcription regulators. RNF 10 (RING finger protein 10), a 90-kDa protein, contains a ring finger motif which is known to be involved in protein-protein interactions. RNF 10 regulates myelin-associated glycoprotein (MAG) expression and is required for myelin formation, therefore, participates in the peripheral nerve development and Schawnn cell differentiation. RNF 10 is also involved in activation of p21WAF 1 promoter via its activity as transcriptional factor.

Notable Publications

Author	Pubmed ID	Journal	Application
Nicolò Carrano	31069631	Mol Neurobiol	WB,IHC
Margarita C Dinamarca	26977767	Elife	WB,IF
Aitor Garzia	34348161	Cell Rep	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



PC-3 cells were subjected to SDS PAGE followed by western blot with 16936-1-AP (RNF10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours




Various lysates were subjected to SDS PAGE followed by western blot with 16936-1-AP (RNF10 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




IP result of anti-RNF10 (IP:16936-1-AP, 4ug; Detection:16936-1-AP 1:500) with PC-3 cells lysate 1480 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16936-1-AP (RNF10 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 16936-1-AP (RNF 10 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RNF10 antibody (16936-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).