For Research Use Only

CHAF1A Polyclonal antibody

Catalog Number: 17037-1-AP

Featured Product

7 Publications



Purification Method:

WB 1:1000-1:4000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 17037-1-AP BC067093 GeneID (NCBI): Size: $600 \mu g/ml$ 10036 **UNIPROT ID:** Source: Rabbit Q13111 Full Name:

chromatin assembly factor 1, subunit

A (p150) Immunogen Catalog Number:

AG10536 Calculated MW:

956 aa, 107 kDa Observed MW: 150 kDa

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IF, IHC, ChIP Species Specificity: human, mouse

Cited Species: human, mouse

Isotype:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, HeLa cells

IP: Jurkat cells,

IHC: human cervix tissue,

Background Information

Chromatin assembly factor 1 (CAF1) is the only histone chaperone known to assemble histones H3 and H4 onto newly synthesized DNA both in vitro and in vivo [PMID:17065558]. The 938 amino acid multidomain p150 (CHAF1A) binds via its C-terminal third to p60, which is an essential step for nucleosome assembly because knocking down either subunit disrupts the activity [PMID:14519857]. In addition, CAF1 facilitates DNA synthesis depending on the binding of the N-terminal 31 residues of p150 to the proliferating cell nuclear antigen (PCNA), which acts as a $sliding\ clamp\ to\ stimulate\ the\ processivity\ of\ DNA\ polymerase\ [PMID:10648606].\ CHAF1A\ regulates\ the\ formation$ of heterochromatin in mammalian cells during replication and in plants it maintains the transcription of certain subsets of genes. Furthermore, CHAF1A exists in a chromatin-remodeling complex WINAC, which coactivates ligand-induced transactivation function of the vitamin D receptor [PMID:12837248]. CHAF1A protein exists some phosphorylation sites, which may affect its theoretical molecular weight when tested. And a 150 kDa band was recognized (PMID:27445493).

Notable Publications

Author	Pubmed ID	Journal	Application
Nasim A Begum	22843687	J Biol Chem	WB
Takashi Ishiuchi	33169018	Nat Struct Mol Biol	WB
Xiancai Ma	33739466	EMBO J	ChIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 17037-1-AP (CHAF1A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-CHAF1A (IP:17037-1-AP, 4ug; Detection:17037-1-AP 1:1000) with Jurkat cells lysate 2400ug.



Immunohistochemical analysis of paraffinembedded human cervix tissue slide using 17037-1-AP (CHAF1A Antibody) at dilution of 1:50 (under 40x lens).