

# METAP2 Polyclonal antibody

Catalog Number: 17040-1-AP

1 Publications

## Basic Information

## Catalog Number:

17040-1-AP

## Size:

700 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG10564

## GenBank Accession Number:

BC013782

## GeneID (NCBI):

10988

## UNIPROT ID:

P50579

## Full Name:

methionyl aminopeptidase 2

## Calculated MW:

478 aa, 53 kDa

## Observed MW:

67 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF/ICC 1:50-1:500

## Applications

## Tested Applications:

IF/ICC, IP, WB, ELISA

## Cited Applications:

WB

## Species Specificity:

human, mouse, rat

## Cited Species:

mouse

## Positive Controls:

WB : HepG2 cells, MCF-7 cells, mouse pancreas tissue, mouse testis tissue

IP : MCF-7 cells,

IF/ICC : HepG2 cells,

## Background Information

MetAP2, also known as MAP2, P67EIF2 or p67, is one of the three known methionine aminopeptidases responsible for the removal of the N-terminal translation initiator methionine from newly synthesized proteins, which is a critical step in protein maturation (PMID: 27692546, 16568094). The enzyme binds two cobalt or manganese ions in its active site (PMID: 11524009, 12718546). As higher MetAP2 expression was observed in tumor (mesothelioma, neuroblastoma, and colorectal carcinoma) cells compared with normal cells, the enzyme serves as a target for anti-angiogenic compounds of natural origin, such as fumagillin and ovalicin (PMID: 11485930, 15927358, 15102683, 9224570). Selective inhibition of MetAP2 stops vascularization and tumor growth in animal models (PMID: 12118091). MetAP2 has 3 isoforms with molecular mass of 50-53 kDa and apparent molecular mass 67 kD.

## Notable Publications

Author	Pubmed ID	Journal	Application
Kotaro Fujii	30576652	Mol Cell	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17040-1-AP (METAP2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



IP result of anti-METAP2 (IP:17040-1-AP, 4ug; Detection:17040-1-AP 1:300) with MCF-7 cells lysate 3200ug.



Immunofluorescent analysis of (-20℃ Ethanol) fixed HepG2 cells using 17040-1-AP (METAP2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).