For Research Use Only

TBC1D5 Polyclonal antibody

Catalog Number: 17078-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17078-1-AP BC013145

Concentration: GeneID (NCBI): 9779

Source: UNIPROT ID: Rabbit Q92609

Isotype: Full Name:

IgG TBC1 domain family, member 5

Immunogen Catalog Number:Calculated MW:AG10991795 aa, 89 kDaObserved MW:

Observed 89 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500

Applications

Tested Applications: WB, IP, IHC, ELISA Cited Applications: WB. IF

Species Specificity: human, mouse Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HEK-293T cells, A431 cells, Jurkat cells, PC-3 cells

IP: A431 cells,

IHC: mouse testis tissue,

Background Information

TBC1D5, a member of TBC (Tre2/Bub2/Cdc16)1 domain family, is a novel retromer-interacting protein. TBC1D5 is a Rab GTPase-activating proteins (GAPs) proteins that negatively regulates VPS35/29/26 recruitment and causes Rab7 to dissociate from the membrane. TBC1D5 could bridge the endosome and autophagosome via its C-terminal LIR motif, and Rab GAPs are implicated in the reprogramming of the endocytic trafficking events under starvation-induced autophagy. The TBC1D5 protein exists 89 kDa and 91 kDa isoforms.

Notable Publications

Author	Pubmed ID	Journal	Application
Ana Jimenez-Orgaz	29158324	EMBO J	WB
Diyuan Zhou	35637944	Int J Biol Sci	WB
Kai-En Chen	34851660	Sci Adv	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17078-1-AP (TBC 1D5 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 17078-1-AP (TBC 1D5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 17078-1-AP (TBC 1D5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 17078-1-AP (TBC 1D5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-TBC 1D5 (IP:17078-1-AP, 4ug; Detection:17078-1-AP 1:300) with A431 cells lysate 2400ug.