For Research Use Only

DGAT2 Polyclonal antibody

Catalog Number:17100-1-AP 5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17100-1-AP BC015234 GeneID (NCBI): Size: 600 ug/ml 84649 **UNIPROT ID:** Source: Rabbit Q96PD7 Full Name: Isotype:

diacylglycerol O-acyltransferase

homolog 2 (mouse)

Immunogen Catalog Number: AG10719 Calculated MW: 388 aa, 44 kDa Observed MW:

39~44kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IF

Species Specificity: human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse adipose tissue, LO2 cells, MCF-7 cells, Neuro-2a cells, mouse heart tissue, mouse liver tissue

Purification Method:

WB 1:500-1:1000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human liver tissue.

Background Information

Diacylglycerol acyltransferase 2 (DGAT2) is an integral membrane protein and major enzyme that catalyzes the final step of triacylglycerol (TG) synthesis in eukaryotes (PMID: 21680734). DGAT2 is present in mitochondriaassociated membranes, specialized domains of the ER that are highly enriched in lipid synthetic enzymes (PMID: 19049983). DGAT2 isoforms (39 kDa, 44kDa) have major differences in acyl-CoA substrate specificity toward erucoyl-CoA.

Notable Publications

Author	Pubmed ID	Journal	Application
Huiduo Guo	34695613	Ecotoxicol Environ Saf	WB
Wei Chen	39549860	Biochim Biophys Acta Mol Cell Biol Lipids WB	
Sami T Tuomivaara	39210032	Commun Biol	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17100-1-AP (DGAT2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 17100-1-AP (DGAT2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 17100-1-AP (DGAT2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.