For Research Use Only

UPF3A Polyclonal antibody

Catalog Number:17114-1-AP

Featured Product





Basic Information

17114-1-AP Size: 300 μ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10799

Catalog Number:

GenBank Accession Number: BC023569 GeneID (NCBI): 65110 UNIPROT ID: Q9H1J1 Full Name: UPF3 regulator of nonsense transcripts homolog A (yeast) Calculated MW: 443 aa, 51 kDa Observed MW:

Positive Controls:

IF/ICC : HeLa cells,

Jurkat cells

55 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:20-1:200

WB: HEK-293 cells, HeLa cells, human testis tissue,

IHC : human testis tissue, human heart tissue

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC Species Specificity: human Cited Species: human Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; ()* Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Notable Publications Author Pubmed ID Journal Application Xinmin Bao 32606924 Cancer Manag Res WB **Dairong Feng** 25934542 **Biochim Biophys Acta** WB 39768178 Cells WB Sara Manore

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Immunohistochemical analysis of paraffinembedded human testis tissue slide using 17114-1-AP (UPF3A Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 17114-1-AP (UPF3A Antibody) at dilution of 1:200 (under 40x lens).

Various lysates were subjected to SDS PAGE followed by western blot with 17114-1-AP (UPF3A antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HeLa cells using 17114-1-AP (UPF3A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated Goat Anti-Rabbit IgG(H+L).