For Research Use Only

METTL11A Polyclonal antibody

Catalog Number:17184-1-AP




Basic Information	Catalog Number: 17184-1-AP	GenBank Accession Number: BC033234	Purification Method: Antigen affinity purification	
	Size: 550 ug/ml Source: Rabbit	GeneID (NCBI): 28989 UNIPROT ID: Q9BV86	Recommended Dilutions: WB 1:1000-1:8000 IHC 1:50-1:500	
				lsotype: IgG
	Immunogen Catalog Number: AG10967	Calculated MW: 223 aa, 25 kDa Observed MW: 16 kDa, 26 kDa		
	Applications	Tested Applications: WB, IHC, ELISA	Positiv	Positive Controls:
Species Specificity: testis		431 cells, HEK-293T cells, HepG2 cells, mouse tissue, rat testis tissue		
human, mouse, rat		IHC : human ovary cancer tissue,		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	METTL11A, also known as NTMT1 (N a -trimethyllysine hydrolase), is an enzyme that belongs to the methyltransferase-like (METTL) protein family. It plays a crucial role in epigenetic processes, particularly in the methylation of proteins, which can influence gene expression and other cellular functions. METTL11A is regulated by other members of the METTL family, such as METTL11B and METTL13, which can either activate or inhibit its function.			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	1 50% glycerol pH 7.3.		

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 17184-1-AP (METTL11A antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 17184-1-AP (METTL11A antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.