
For Research Use Only

G-CSF Polyclonal antibody

Catalog Number:17185-1-AP

Featured Product







Basic Information	Catalog Number: 17185-1-AP	GenBank Accession Number: BC033245	Purification Method: Antigen affinity purification	
	Size: 550 µg/ml	GeneID (NCBI): 1440	Recommended Dilutions: WB 1:500-1:2000	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10968	UNIPROT ID: P09919 Full Name: colony stimulating factor 3 (granulocyte)		
				Calculated MW: 200 aa, 22 kDa
		Observed MW: 20 kDa		
		Applications	Tested Applications:	Positi
WB, ELISA, FC (Intra) Cited Applications: WB, IHC				
Species Specificity: human				
Cited Species: human				
Background Information	Granulocyte colony-stimulating factor (G-CSF), also referred to as CSF3, is a protective cytokine with anti- inflammatory effects. G-CSF is important in promoting survival of the granulocytic lineage cells and proliferation and migration of neutrophils as well as trophoblast cells. G-CSF acts by binding to its receptor G-CSFR (also called CSF3R), which after binding with G-CSF activates the canonical Janus kinase (Jak)/signal transducer, activator of transcription (STAT)and Ras/Raf/MAP kinase pathways. G-CSF potently stimulates the proliferation and release of peripheral blood progenitor cells into the bloodstream and is therefore used to treat neutropenia after chemotherapy. Furthermore, G-CSF levels are elevated upon intensive exercise leading to increased neutrophil counts, which are predominantly due to delayed neutrophil apoptosis.			
Notable Publications	Author	Pubmed ID Journal	Application	
	Zheng Zhang	31765840 Mol Immunol		
	Wei Jiang	36982170 Int J Mol Sci	WB,IHC	
Storage	Storage: Store at -20°C. Stable for one yes Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -21	nd 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 17185-1-AP (G-CSF antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

1X10^6 K-562 cells were intracellularly stained with 0.4 ug Anti-Human G-CSF (17185-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).