

For Research Use Only

BBX Polyclonal antibody

Catalog Number: 17254-1-AP

Featured Product

4 Publications



Basic Information

Catalog Number:

17254-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10691

GenBank Accession Number:

BC012837

GeneID (NCBI):

56987

UNIPROT ID:

Q8WY36

Full Name:

bobby sox homolog (Drosophila)

Calculated MW:

941 aa, 105 kDa

Observed MW:

105 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

WB, IP, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

mouse, rat, zebrafish

Positive Controls:

WB : HepG2 cells,

IP : HepG2 cells,

IHC : human ovary tissue, human brain tissue, human lung tissue, human placenta tissue, human spleen tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

BBX, also named HBP2, is a protein with a high mobility group (HMG) domain that belongs to the Sox (Sry-related HMG box) family of transcription factors. As a DNA-binding domain in HBP2, HMG is essential for its function in Schizosaccharomyces pombe. HBP2 activates Cdc10-dependent transcription at G1/s and allows growth of cdc10-129, res1delta and rep2delta mutant cells at the restrictive temperature

Notable Publications

Author	Pubmed ID	Journal	Application
Chieh-Ying Wang	27465138	Biol Reprod	WB, IHC, IF
TieLin Chen	25079045	Dev Genes Evol	IHC
Su Jeong Lee	39482539	Exp Mol Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 17254-1-AP (BBX antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary using 17254-1-AP (BBX antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human ovary using 17254-1-AP (BBX antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-BBX (IP:17254-1-AP, 4ug; Detection:17254-1-AP 1:1000) with HepG2 cells lysate 2400ug.