For Research Use Only

SLC25A18 Polyclonal antibody

Catalog Number: 17348-1-AP 3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17348-1-AP BC031644

Size: GeneID (NCBI): 353 µg/ml 83733

Source: UNIPROT ID: Rabbit Q9H1K4

Isotype: Full Name:

Isotype: Full Name:
IgG solute carrier family 25

Immunogen Catalog Number: (mitochondrial carrier), member 18

Calculated MW:

32-35 kDa

315 aa, 34 kDa Observed MW: Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500

Applications

Tested Applications: WB, IP, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse

Cited Species: human, mouse

AG11420

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, mouse kidney tissue, mouse

heart tissue

IP: mouse brain tissue,
IHC: mouse brain tissue,

Background Information

SLC25A18, also named as GC 2, belongs to the mitochondrial transporter family (SLC25). It is located in inner mitochondrial membrane which involved in the transport of glutamate either with H+ or in exchange for OH-. SLC25A18 is highly expressed in brain, testis, lung, kidney, liver, spleen and skeletal muscle. 17348-1-AP antibody detects the band around 32-35 kDa in SDS-PAGE. (PMID: 14598172)

Notable Publications

Author	Pubmed ID	Journal	Application
Li Liang	32509397	Am J Cancer Res	IHC
Jinghui Liang	39355508	MedComm (2020)	WB
Jia-Yi Song	38871013	Mitochondrion	WB,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 17348-1-AP (SLC25A18 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-SLC 25A18 (IP:17348-1-AP, 4ug; Detection:17348-1-AP 1:500) with mouse brain tissue lysate 4400ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 17348-1-AP (SLC25A18 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 17348-1-AP (SLC25A18 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).