For Research Use Only

MKNK2 Polyclonal antibody Catalog Number: 17354-1-AP Featured Product 5

Featured Product 5 Publications





Basic Information	Catalog Number: 17354-1-AP	GenBank Accession Number: BC018345	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	600 ug/ml	2872	WB 1:200-1:1000	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10546	UNIPROT ID: Q9HBH9	IHC 1:100-1:400 IF/ICC 1:50-1:500	
		Full Name:		
		MAP kinase interacting serine/threonine kinase 2		
		Calculated MW: 465 aa, 52 kDa		
		Observed MW: 47-52 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA	WB : mouse spleen tissue, MCF-7 cells		
	Cited Applications: WB, IHC	IHC : hum	an breast cancer tissue,	
	Species Specificity: human, mouse, rat	IF/ICC : MCF-7 cells,		
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	buffer pH 6.0			
Background Information	MKNK2(MAP kinase-interacting se CAMK Ser/Thr protein kinase fam homology to the calcium/calmod located within the T loop of the ki uncharacterized form of MNK2 or a	ily. It has a conserved C-terminal Erk	interacting domain, a catalytic domain wi d putative MAP kinase phosphorylation site ast cancer cells express either an th the band of 72-75 kDa by western	
	MKNK2(MAP kinase-interacting se CAMK Ser/Thr protein kinase fam homology to the calcium/calmod located within the T loop of the ki uncharacterized form of MNK2 or a blot(PMID:17130135). MKNK2 has	ily. It has a conserved C-terminal Erk- ulin-dependent family of kinases, and nase domain(PMID:9155017). The bre a cross-reacting MNK2-like protein win 2 isoforms with the molecular mass of	th the band of 72-75 kDa by western If 52 kDa and 47 kDa.	
	MKNK2(MAP kinase-interacting se CAMK Ser/Thr protein kinase fam homology to the calcium/calmod located within the T loop of the ki uncharacterized form of MNK2 or a blot(PMID:17130135). MKNK2 has	ily. It has a conserved C-terminal Erk- ulin-dependent family of kinases, an nase domain(PMID:9155017). The bre a cross-reacting MNK2-like protein wi 2 isoforms with the molecular mass of Pubmed ID Journal	interacting domain, a catalytic domain wi d putative MAP kinase phosphorylation site ast cancer cells express either an th the band of 72-75 kDa by western	
	MKNK2(MAP kinase-interacting se CAMK Ser/Thr protein kinase fam homology to the calcium/calmod located within the T loop of the ki uncharacterized form of MNK2 or a blot(PMID:17130135). MKNK2 has	ily. It has a conserved C-terminal Erk- ulin-dependent family of kinases, and nase domain(PMID:9155017). The bre a cross-reacting MNK2-like protein win 2 isoforms with the molecular mass of	interacting domain, a catalytic domain wi d putative MAP kinase phosphorylation site ast cancer cells express either an the band of 72-75 kDa by western of 52 kDa and 47 kDa. Application	
Background Information Notable Publications	MKNK2(MAP kinase-interacting se CAMK Ser/Thr protein kinase fam homology to the calcium/calmod located within the T loop of the ki uncharacterized form of MNK2 or a blot(PMID:17130135). MKNK2 has Author Milagros Suarez	ily. It has a conserved C-terminal Erk- ulin-dependent family of kinases, and nase domain(PMID:9155017). The bre a cross-reacting MNK2-like protein wit 2 isoforms with the molecular mass of Pubmed ID Journal 34012509 Oncotarget	interacting domain, a catalytic domain with d putative MAP kinase phosphorylation site east cancer cells express either an the band of 72-75 kDa by western of 52 kDa and 47 kDa. Application WB	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data








Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 17354-1-AP (MKNK2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 17354-1-AP (MKNK2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using MKNK2 antibody (17354-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).