

For Research Use Only

Plasminogen Polyclonal antibody

Catalog Number: 17462-1-AP

1 Publications



Basic Information

Catalog Number:

17462-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11217

GenBank Accession Number:

BC060513

GeneID (NCBI):

5340

UNIPROT ID:

P00747

Full Name:

plasminogen

Calculated MW:

810 aa, 91 kDa

Observed MW:

97 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, WB, ELISA

Species Specificity:

human

Positive Controls:

WB : human blood tissue, human placenta tissue

IF : HepG2 cells,

Background Information

Plasminogen is a secreted blood zymogen that is activated by proteolysis and converted to plasmin and angiotatin. Plasminogen plays an important role in fibrinolysis as well as wound healing, cell migration, tissue modeling and angiogenesis. Plasminogen is synthesized in the liver, and it circulates in the blood, with a half-life of 2.2 days. Plasminogen is the precursor of plasmin, which lyses fibrin clots to fibrin degradation products (FDP) and D-dimer; the conversion to active protease is mediated by tissue-type (tPA) and urokinase-type (uPA) plasminogen activators. Defects in plasminogen are likely a cause of thrombophilia and ligneous conjunctivitis.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Zhou	34305859	Front Microbiol	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human blood were subjected to SDS PAGE followed by western blot with 17462-1-AP (Plasminogen antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 17462-1-AP (Plasminogen antibody) at dilution of 1:50 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).