

For Research Use Only

Plasminogen Polyclonal antibody

Catalog Number: 17462-1-AP **3 Publications**



Basic Information

Catalog Number: 17462-1-AP	GenBank Accession Number: BC060513	Purification Method: Antigen affinity purification
Size: 500 ug/ml	GeneID (NCBI): 5340	Recommended Dilutions: WB 1:1000-1:4000
Source: Rabbit	UNIPROT ID: P00747	IHC 1:50-1:500 IF/ICC 1:50-1:500
Isotype: IgG	Full Name: plasminogen	
Immunogen Catalog Number: AG11217	Calculated MW: 810 aa, 91 kDa	
	Observed MW: 97 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : human blood tissue, human placenta tissue
Cited Applications: WB	IHC : human stomach tissue,
Species Specificity: human	IF/ICC : HepG2 cells,
Cited Species: mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Plasminogen is a secreted blood zymogen that is activated by proteolysis and converted to plasmin and angiostatin. Plasminogen plays an important role in fibrinolysis as well as wound healing, cell migration, tissue modeling and angiogenesis. Plasminogen is synthesized in the liver, and it circulates in the blood, with a half-life of 2.2 days. Plasminogen is the precursor of plasmin, which lyses fibrin clots to fibrin degradation products (FDP) and D-dimer; the conversion to active protease is mediated by tissue-type (tPA) and urokinase-type (uPA) plasminogen activators. Defects in plasminogen are likely a cause of thrombophilia and ligneous conjunctivitis.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Zhou	34305859	Front Microbiol	
Jonathan J Molina	38538626	Nat Commun	WB
Ivan Ivanov	30591525	Blood	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



human blood were subjected to SDS PAGE followed by western blot with 17462-1-AP (Plasminogen antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 17462-1-AP (Plasminogen antibody) at dilution of 1:50 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human stomach tissue slide using 17462-1-AP (Plasminogen antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).