

For Research Use Only

HEMGN Polyclonal antibody

Catalog Number: 17616-1-AP

1 Publications



Basic Information

Catalog Number:

17616-1-AP

Size:

240 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11793

GenBank Accession Number:

BC048324

GeneID (NCBI):

55363

UNIPROT ID:

Q9BXL5

Full Name:

hemogen

Calculated MW:

483 aa, 55 kDa

Observed MW:

60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

IHC

Species Specificity:

human, mouse

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : Jurkat cells, HL-60 cells, human testis tissue, mouse testis tissue

IP : HL-60 cells,

IHC : human testis tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Yunshu Zhu	37760841	Biomedicines	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 17616-1-AP (HEMGN antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human testis using 17616-1-AP (HEMGN antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human testis using 17616-1-AP (HEMGN antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-HEMGN (IP:17616-1-AP, 4ug; Detection:17616-1-AP 1:500) with HL-60 cells lysate 3000ug.