For Research Use Only

MON1B Polyclonal antibody

Catalog Number: 17638-1-AP

Featured Product

1 Publications





Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 17638-1-AP BC024277 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 650 µg/ml 22879 WB 1:1000-1:4000 IF/ICC 1:50-1:500 UNIPROT ID: Source: Rabbit Q7L1V2 Full Name: Isotype: lgG MON1 homolog B (yeast) Calculated MW: Immunogen Catalog Number: AG9920 547 aa, 59 kDa **Observed MW:** 59 kDa **Applications Tested Applications:** Positive Controls: WB, IF/ICC, ELISA WB : HeLa cells, Jurkat cells, K-562 cells **Cited Applications:** IF/ICC : HeLa cells, WB, IF Species Specificity: human **Cited Species:** human **Background Information** Vacuolar fusion protein MON1 homolog B (MON1B) is also named as HSRG1, KIAA0872 and SAND2, and belongs to the MON1/SAND family. The membrane recruitment of MON1B, a homolog of yeast MON1 protein in mammals, is essential for vesicular fusion of early endosomes (PMID:26987402). Notable Publications Author Pubmed ID Application Journal Ximing Shao 26987402 Cell Res WB,IF Storage Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





WB result of MON1B antibody (17638-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MON1B transfected HeLa cells.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using MON1B antibody (17638-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 17638-1-AP (MON1B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.