

For Research Use Only

CD226 Polyclonal antibody

Catalog Number: 17842-1-AP **1 Publications**



Basic Information

Catalog Number: 17842-1-AP	GenBank Accession Number: BC074787	Purification Method: Antigen affinity purification
Size: 550 µg/ml	GeneID (NCBI): 10666	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	ENSEMBL Gene ID: ENSG00000150637	
Isotype: IgG	UNIPROT ID: Q15762	
Immunogen Catalog Number: AG12206	Full Name: CD226 molecule	
	Calculated MW: 336 aa, 39 kDa	
	Observed MW: 65-70 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : Human peripheral blood platelets, Jurkat cells IHC : human tonsillitis tissue,
Cited Applications: WB	
Species Specificity: human	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

CD226 (DNAM-1) is a ~65 kDa glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. CD226 mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Interactions of CD226 and its ligands, CD155 and CD112, induce NK and T cell-mediated cytotoxicity and cytokine secretion (PMID: 15039383).

Notable Publications

Author	Pubmed ID	Journal	Application
Yidong Chen	37083227	Adv Sci (Weinh)	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 17842-1-AP (CD226 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17842-1-AP (CD226 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 17842-1-AP (CD226 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).