For Research Use Only

SFRS9 Polyclonal antibody

Catalog Number: 17926-1-AP

Featured Product

7 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17926-1-AP BC093973
Size: GeneID (NCBI): 450 µg/ml 8683
Source: UNIPROT ID: Rabbit Q13242

Recommended Dilutions: WB 1:500-1:2000 IHC 1:500-1:2000

Purification Method:

Antigen affinity purification

Isotype: Full Name:

IgG splicing factor, arginine/serine-rich 9

Immunogen Catalog Number:Calculated MW:AG12347221 aa, 26 kDaObserved MW:

26 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells
IHC: mouse kidney tissue, human cervical cancer
tissue, human urothelial carcinoma tissue

Background Information

SFRS9 is a member of the serine/arginine (SR) splicing factor family, which contains an N-terminal RNA recognition motif (RRM), a glycine-rich region, an internal region homologous to the RRM, and a long (315-amino-acid) C-terminal domain composed predominantly of alternating serine and arginine residues. SRSF9 has been linked to the occurrence and progression of hepatocellular carcinoma (HCC), colorectal cancer (CRC) and oral squamous cell carcinoma (OSCC). In HCC and CRC, the knockdown of SRSF9 inhibited the proliferation and migration of cancer cells, cell cycle progression and colony formation of cancer cells (PMID: 34336668,35971121).

Notable Publications

Author	Pubmed ID	Journal	Application
Chi-Ren Tsai	30260058	Hum Mutat	WB
Haiyan An	31636118	J Cell Biol	WB
Haiyang Zhang	34845198	Cell Death Dis	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 17926-1-AP (SFRS9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 17926-1-AP (SFRS9 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 17926-1-AP (SFRS9 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).