

For Research Use Only

Neurokinin-1 receptor Polyclonal antibody



Catalog Number: 17942-1-AP

6 Publications

Basic Information

Catalog Number:

17942-1-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12368

GenBank Accession Number:

BC074911

GeneID (NCBI):

6869

UNIPROT ID:

P25103

Full Name:

tachykinin receptor 1

Calculated MW:

407 aa, 46 kDa

Observed MW:

45-55 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : SH-SY5Y cells, THP-1 cells, mouse pancreas tissue, Caco-2 cells, HeLa cells

Background Information

neurokinin 1 receptor (NK1R, also known as the tachykinin 1 receptor, TACR1), a mediator of behavioral stress responses, in alcohol dependence and treatment. One such neurotransmitter is substance P (SP), which together with its preferred TACR1 is highly expressed in brain areas involved in stress responses and drug reward, including the hypothalamus, amygdala, and nucleus accumbens (PMID: 18276852).

Notable Publications

Author	Pubmed ID	Journal	Application
Ping Ma	34710756	Phytomedicine	WB
Yan Yu	29759144	J Neuroimmunol	WB
Xiao-Dan Guo	32853022	Neoplasma	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17942-1-AP (Neurokinin-1 receptor antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 17942-1-AP (Neurokinin-1 receptor antibody at dilution of 1:600 incubated at room temperature for 1.5 hours.