For Research Use Only

RGN/SMP30 Polyclonal antibody

Catalog Number:17947-1-AP 11 Publications



Basic Information

Catalog Number: GenBank Accession Number: 17947-1-AP BC073173 GeneID (NCBI): Size: 400 μg/ml 9104 **UNIPROT ID:** Source: Rabbit Q15493 Full Name: Isotype:

regucalcin (senescence marker protein-30)

AG12423 Calculated MW: 299 aa. 33 kDa

Observed MW: 33 kDa. 25 kDa **Purification Method:** Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

WB 1:3000-1:10000

Applications

Tested Applications: WB, IP, IHC, ELISA **Cited Applications:** WB, IHC, IF Species Specificity: human, mouse, rat **Cited Species:**

Immunogen Catalog Number:

human, mouse, rat, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse liver tissue, mouse small intestine tissue,

rat liver tissue

IP: mouse liver tissue.

IHC: rat kidney tissue, human liver tissue, human kidney tissue, human hepatocirrhosis tissue

Background Information

 $Regucal cin \ (RGN), also \ known \ as \ SMP30, is \ a \ highly \ conserved, calcium-binding \ protein, that \ is \ preferentially$ expressed in the liver and kidney. SMP30 is preferentially localized in hepatocytes and renal proximal tubular epithelial cells. It is known to be related to aging, hepatocyte proliferation and tumorigenesis. It may be useful for HCC serologic screening, especially for the patients with AFP Negative. RGN encodes two isoforms: 33 kDa and 25

Notable Publications

Author	Pubmed ID	Journal	Application
Le Zhang	34688152	Int Immunopharmacol	WB
Norihiro Okada	34082065	Gene	WB
Xiaolong Li	31289533	Oncol Lett	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 17947-1-AP (RGN/SMP30 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP result of anti-RGN/SMP30 (IP:17947-1-AP, 4ug; Detection:17947-1-AP 1:15000) with mouse liver tissue lysate 3040 ug.



Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 17947-1-AP (RGN/SMP30 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat kidney tissue slide using 17947-1-AP (RGN/SMP30 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).