

For Research Use Only

EDAR Polyclonal antibody

Catalog Number: 18032-1-AP

4 Publications



Basic Information

Catalog Number:

18032-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG12601

GenBank Accession Number:

BC093872

GeneID (NCBI):

10913

UNIPROT ID:

Q9UNEO

Full Name:

ectodysplasin A receptor

Calculated MW:

448 aa, 49 kDa

Observed MW:

48 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2400

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications:

WB, IP, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human

Cited Species:

mouse

Positive Controls:

WB : A549 cells, HEK-293 cells, A431 cells, MCF-7 cells

IP : A549 cells,

Background Information

EDAR, also named as DL, is a receptor for EDA isoform A1, but not for EDA isoform A2. It mediates the activation of NF-kappa-B and JNK. EDAR may promote caspase-independent cell death. It is expressed in adjacent or partially overlapping layers in the developing human skin with MW 48 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Chang-Yi Cui	25249463	Development	IF
Zhong-San Zeng	30663067	J Pharm Pharmacol	WB
Qirui Wang	38582368	J Invest Dermatol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



A549 cells were subjected to SDS PAGE followed by western blot with 18032-1-AP (EDAR antibody) at dilution of 1:1200 incubated at room temperature for 1.5 hours.



IP result of anti-EDAR (IP:18032-1-AP, 4ug; Detection:18032-1-AP 1:500) with A549 cells lysate 2800ug.



Various lysates were subjected to SDS PAGE followed by western blot with 18032-1-AP (EDAR antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.