## For Research Use Only

## CYP7A1 Polyclonal antibody, PBS Only

Catalog Number: 18054-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** 

18054-1-PBS

BC101777 GeneID (NCBI): Antigen affinity purification

Size: 1 mg/ml Source:

1581 **UNIPROT ID:** P22680

Full Name:

Rabbit Isotype:

cytochrome P450, family 7, subfamily

Immunogen Catalog Number: AG12572

A, polypeptide 1 Calculated MW:

504 aa, 58 kDa

Observed MW: 58 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

**Background Information** 

CYP7A1 (Cytochrome P450 Family 7 Subfamily A Member 1), is a cholesterol 7alpha-hydroxylase, catalyzing the first and rate-limiting step in the neutral or classic pathway for bile acid biosynthesis. CYP7A1 is a postulated gene modifier of colorectal cancer risk and target for the therapeutic bile acid, ursodeoxycholic acid (UDCA). The CYP7A1 enzyme controls the rate-limiting step of the catabolism of cholesterol into bile acids in the liver.


Storage

Storage:


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 18054-1-AP (CYP7A1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 18054-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 18054-1-AP (CYP7A1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 18054-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 18054-1-AP (CYP7A1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 18054-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CYP7A1 antibody (18054-1-AP) at dilution of 1:200 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). This data was developed using the same antibody clone with 18054-1-PBS in a different storage buffer formulation.